

Clinical response to the NEWS2 trigger thresholds

NEW score	Frequency of monitoring	Clinical response
0	Minimum 12 hourly	<ul style="list-style-type: none"> Continue routine NEWS monitoring
Total 1–4	Minimum 4–6 hourly	<ul style="list-style-type: none"> Inform registered nurse, who must assess the patient Registered nurse decides whether increased frequency of monitoring and/or escalation of care is required
3 in single parameter	Minimum 1 hourly	<ul style="list-style-type: none"> Registered nurse to inform medical team caring for the patient, who will review and decide whether escalation of care is necessary
Total 5 or more Urgent response threshold	Minimum 1 hourly	<ul style="list-style-type: none"> Registered nurse to immediately inform the medical team caring for the patient Registered nurse to request urgent assessment by a clinician or team with core competencies in the care of acutely ill patients Provide clinical care in an environment with monitoring facilities
Total 7 or more Emergency response threshold	Continuous monitoring of vital signs	<ul style="list-style-type: none"> Registered nurse to immediately inform the medical team caring for the patient – this should be at least at specialist registrar level Emergency assessment by a team with critical care competencies, including practitioner(s) with advanced airway management skills Consider transfer of care to a level 2 or 3 clinical care facility, ie higher-dependency unit or ICU Clinical care in an environment with monitoring facilities